

SHORT COMMUNICATIONS

J. Raptor Res. 50(2):211–216

© 2016 The Raptor Research Foundation, Inc.

THE IMMINENT DISAPPEARANCE OF THE APLOMADO FALCON FROM THE CHIHUAHUAN DESERT

ALBERTO MACÍAS-DUARTE¹

*Universidad Estatal de Sonora, Cuerpo Académico de Recursos Naturales, Unidad Académica Hermosillo,
Ley Federal del Trabajo S/N, Col. Apolo, Hermosillo, Sonora 83100 Mexico*

ANGEL B. MONTOYA²

The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709 U.S.A.

J. ROBERTO RODRÍGUEZ-SALAZAR

The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709 U.S.A.

and

*IMC Vida Silvestre A.C., Río Jordán 4311, Col. Junta de los Ríos, Chihuahua,
Chihuahua 31300 Mexico*

ARVIND O. PANJABI

Bird Conservancy of the Rockies, 230 Cherry, Suite 130, Fort Collins, CO 80521 U.S.A.

PEDRO A. CALDERÓN-DOMÍNGUEZ

IMC Vida Silvestre A.C., Río Jordán 4311, Col. Junta de los Ríos, Chihuahua, Chihuahua 31300 Mexico

W. GRAINGER HUNT

The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, ID 83709 U.S.A.

KEY WORDS: *Aplomado Falcon*; *Falco femoralis*; *Chihuahuan Desert*; *extirpation*; *Mexico*.

The Northern Aplomado Falcon (*Falco femoralis septentrionalis*) is a savannah-dwelling raptor whose range centers in Latin America, and whose distribution once extended northward into coastal Texas, western Texas, southern New Mexico, and southeastern Arizona (Keddy-Hector 2000, Fig. 1). The species had virtually disappeared from the United States by the 1930s, primarily as a result of habitat destruction by farming and livestock grazing (Hector 1981). A fairly robust population breeds in tropical savannahs from Veracruz, Mexico, southward into Central America, whereas only two small populations occur north of that region. One consists of about 30 pairs restored in recent years to coastal Texas by The Peregrine Fund through captive breeding and release (Hunt et al. 2013). The other occurs in the grasslands of central Chihuahua, Mexico (Macías-Duarte et al. 2004) and is the subject of this paper.

This native population in Chihuahua, discovered in the 1990s, is believed to be the last remnant of the desert-

dwelling Aplomado Falcons that formerly extended into the Chihuahuan Desert region of the United States (Montoya 1995, Montoya et al. 1997, Macías-Duarte et al. 2004). Indeed, there are occasional sightings of wild Aplomado Falcons in nearby southern New Mexico (Young et al. 2004), with several cases of successful nesting just inside its border (Meyer and Williams 2005). Even so, a decade-long attempt at reintroduction of captive-bred falcons in New Mexico and west Texas by The Peregrine Fund produced no self-sustaining population (Hunt et al. 2013). We here report the current status of the wild population in Chihuahua and describe the factors that are undermining its persistence.

Nest-territory monitoring, which began soon after the discovery of the Chihuahua population, revealed a strong dependence of reproductive success and nest initiation date upon annual variation in the abundance of grassland birds (falcon prey), and they, in turn, upon precipitation (Montoya et al. 1997, Macías-Duarte et al. 2004, Hunt et al. 2013). Trends in nesting success suggested that this population was declining by the early 2000s. Overall mean productivity during that period (0.83 fledglings per mated pair per year) appeared too low to sustain a stable population (Macías-Duarte 2002). Droughts, coupled with overgrazing by cattle and falcons drowning in livestock watering tanks, were

¹ Email address: alberto.macias@ues.mx

² Present address: Partners for Fish and Wildlife Program, U.S. Fish and Wildlife Service, 1800 Marquess Street, Las Cruces, NM 88005 U.S.A.




Figure 1. Breeding distribution of the Northern Aplomado Falcon (black areas). The Sueco (west) and Tinaja Verde (east) study areas are shown within the state of Chihuahua in northern Mexico. Map redrawn from Keddy-Hector (2000) and Hunt et al. (2013).

hypothesized as primary causes of population decline. A new threat appeared in 2006 when the creation and development of agricultural colonies began an extensive conversion of occupied falcon habitat into almost 70,000 ha of irrigated farmland (Macías-Duarte et al. 2009, Pool et al. 2014). Besides direct destruction of habitat, farming also brought other potentially adverse factors, including pesticides and an extensive network of roads and electric lines to the region. All things considered, the long-term viability of Aplomado Falcon populations in the Chihuahuan Desert appears increasingly uncertain. Our objective in this report is to document the trend of the Aplomado Falcon population in Chihuahua, under the simultaneous effects of drought and breeding habitat loss and deterioration. We also consider management options for preventing extirpation.

METHODS

We conducted Aplomado Falcon surveys as described in Macías-Duarte et al. (2004) in two study areas, Sueco and Tinaja Verde (Fig. 1) during 2003–2015. The areas differ somewhat in ecological conditions (Macías-Duarte et al. 2004) and the extent of breeding habitat destruction. Falcon habitat at Sueco is hotter, drier, flatter, and lower in elevation than that at Tinaja Verde; falcon habitat loss by farming has occurred only at Sueco.

We defined a breeding territory (territory hereafter) as an area that contained at least one nest from 1997–2014 within the range of a mated pair. We defined an occupied territory as one containing a mated pair, an active territory as one where eggs were laid, and a successful territory as one containing large young. Our approach in monitoring falcon reproduction was to find and monitor as many occupied and active territories as possible within the two study areas. By 2002, Macías-Duarte et al. (2004) had identified and monitored 24 territories at Sueco and 11 at Tinaja Verde, and we

did not continue searching for more in this study. We defined an “extant” territory as one occupied by a pair at least once since 2000 and still intact; i.e., not yet converted to cropland. We began each annual survey prior to incubation and periodically observed pairs until fledgling or failure was confirmed. Prior to 2012, we visited each occupied and previously occupied territory at least twice per month from February to June. We surveyed only once per month from 2012–2014. In some cases, territories noted as unoccupied may have contained undetected pairs. In 2015, we surveyed each extant territory once during the entire breeding season, except when we detected a falcon pair, in which case, we monitored it once per month to determine nest success.

RESULTS

The total number of known occupied and active territories steeply declined after 2006 in both Sueco and Tinaja Verde (Fig. 2). The number of occupied territories declined from 31 in 2000 (Macías-Duarte et al. 2004) to three in 2014. Likewise, the number of active territories fell from 24 in 2000 to two in 2014. This decline in territory occupancy was due to both conversion to cropland at Sueco by farming (Fig. 2, 3) and failure of Aplomado Falcons to occupy 19 extant territories, particularly in Tinaja Verde. Conversion of desert grasslands to farmland had not reached the Tinaja Verde study area in 2014 (Fig. 3), and yet we detected only one pair among the 11 territories available there during that breeding season. In 2015, a year of increased precipitation, we found four active territories in Sueco and one in Tinaja Verde, respectively, from which three pairs raised at least two fledglings overall.

DISCUSSION

The observed trends in territory occupancy through the study period suggest a decline, almost to extirpation, of the only known breeding population of Aplomado Falcons in the Chihuahuan Desert. The actual extent of population distribution is unknown because there is no current assessment of the range and quality of open grassland habitat in the region. However, based on trends in monitoring and sightings of pairs and individual falcons outside our study areas, we estimate that there may be as few as 10 pairs remaining in central Chihuahua, which may have considerable consequences for recruitment. Consider, for example, that the 2014 surveys in our study revealed only two nests, collectively producing only one juvenile, as compared with 24 breeding pairs producing 33 juveniles in 2000 (Macías-Duarte 2002). If the factors mediating this decline operate in a regional context—and there is little reason to think otherwise—then the extirpation of the central Chihuahua population appears imminent. In this regard, there was scarce evidence of the presence of Aplomado Falcons in other regions of Chihuahua (although occasional sightings of adults occur outside our study areas), nor in the grasslands of Durango and Coahuila. Habitat protection and enhancement at Sueco and Tinaja Verde are thus important strategies for survival, and the need is urgent.




Figure 2. Trends in occupancy of historical Aplomado Falcon breeding territories and in the numbers of breeding pairs from 1997 to 2014 at Sueco (upper) and Tinaja Verde (lower) study areas in desert grasslands of central Chihuahua, Mexico. Data from 1997–2002 taken from Macías-Duarte et al. (2004). Extant territories were those occupied at least once during the study period and not yet destroyed by farmland conversion.



Figure 4. Aplomado Falcon nest adjacent to farmland in Chihuahua, Mexico (photograph by J.R. Rodríguez-Salazar).

and other management practices are worthy of immediate consideration in view of the predicted negative consequence of inaction.

Given that breeding success is intrinsically linked with summer precipitation, however, climate-change-related drought may nevertheless preclude the recovery of Aplomado Falcons in the Chihuahuan Desert. Ironically, the farmers, by bringing water to the surface for irrigation, might, under the right circumstances, increase prey numbers; e.g., doves, even to the point of sustaining local pairs of Aplomado Falcons. There is little current evidence of such a possibility, however, except in the case of the only successful nest in 2014, a formerly occupied territory adjacent to farmland (Fig. 4). Perhaps not coincidentally, the only other productive site reported in the Chihuahuan Desert in 2014 was just north of the border in New Mexico, ca. 7.2 km from agricultural fields. In addition, two of the five nests found in 2015 were in the vicinity of farmland. It is thus conceivable that, if agricultural practices could maintain chemically safe prey populations year-round, and some of the other measures listed above were implemented, aplomado pairs in Chihuahua might reproduce even in periods of low precipitation. Chemically safe farming in Chihuahua might also promote the survival of juveniles during the post-fledgling dependence period and even nonbreeding

aplomados during drought. A telemetry study of two juveniles in 2015 showed they predominantly used recently opened agricultural areas (Fig. 3), after leaving their natal territories in native grasslands (A. Macías-Duarte unpubl. data). Lastly, the only documented sighting of an aplomado in west Texas in the decades preceding the release program was of an adult frequenting an irrigated field near the town of Valentine during January–May 1992 (Haynie 1994).

DESAPARICIÓN INMINENTE DE *FALCO FEMORALIS* DEL DESIERTO CHIHUAHUENSE

RESUMEN.—Seguimos la ocupación territorial de *Falco femoralis* durante un periodo de 12 años (2003–2014) en el centro de Chihuahua, México, bajo condiciones de sequía severa y destrucción de su hábitat de anidamiento. La conversión de los territorios de cría de los halcones a tierras agrícolas ha destruido hasta ahora 17 de los 35 territorios de cría ocupados por parejas de *F. femoralis* conocidos al menos una vez desde el año 2000. Sólo una pequeña fracción territorio de los restantes fue ocupada en 2014. La desaparición de los halcones de los territorios de cría antiguos que se mantienen intactos sugiere un déficit de reclutamiento causado por tasas reproductivas

bajas relacionadas con una disminución general en la calidad de los pastizales y en la cantidad de presas. La persistencia de unas pocas parejas sugiere que la población se mantiene actualmente por la elevada supervivencia de los adultos más que por su fertilidad. Esta tendencia, así como la continua destrucción del escaso hábitat de alta calidad para *F. femoralis* en Chihuahua y el norte de México, pronostican la desaparición de esta población de *F. femoralis* que habita en el desierto.

[Traducción del equipo editorial]

ACKNOWLEDGMENTS

We thank The Peregrine Fund, T & E Inc., PROFAUNA, World Wildlife Fund, U.S. Fish and Wildlife Service, Canadian Wildlife Service, the Timken Foundation, Bird Conservancy of the Rockies, Professional Development Program for Teachers (PRODEP), Universidad Estatal de Sonora, and IMC Vida Silvestre A.C. for financial support. We thank Alberto Lafón, whose support was instrumental. We thank Secretaría de Medio Ambiente y Recursos Naturales de Mexico for issuing the scientific collection permits that allowed us to work in Mexico (Permits No. SGPA/DGVS/01626, SGPA/DGVS/03546, SGPA/DGVS/07122/09, SGPA/DGVS/02400/13, SGPA/DGVS/09558/13, SGPA/DGVS/01112/15). We especially thank the landowners for providing access to their land, particularly Enrique Baeza and the Borunda Carrillo family who also provided logistical support. Field assistance for this project was provided by Julio Gallardo, Alejandro Nava, Josué Méndez, and Joel Morales. Comments by Tom Cade and two anonymous reviewers greatly improved an earlier version of this report.

LITERATURE CITED

ANONYMOUS. 2010. Norma oficial Mexicana NOM-059-SEMARNAT-2010, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo. Pages 1-77 in Anonymous. Diario Oficial de la Federación, December the 30th (2nd Section). Distrito Federal, Mexico.

HAYNIE, C.B. 1994. Texas bird records committee report for 1994. Texas Ornithology Society annual report. www.texasbirdrecordscommittee.org/home/reports/1994-annual-report (last accessed 28 October 2015).

HECTOR, D.P. 1981. The habitat, diet, and foraging behavior of the Aplomado Falcon, *Falco femoralis* (Temminck). M.S. thesis, Oklahoma State University, Stillwater, OK U.S.A.

HUNT, W.G., J.L. BROWN, T.J. CADE, J. COFFMAN, M. CURTI, E. GOTT, B. HEINRICH, J.P. JENNY, P. JUERGENS, A.

MACÍAS-DUARTE, A.B. MONTOYA, B. MUTCH, AND C. SANDFORT. 2013. Restoring Aplomado Falcons to the United States. *Journal of Raptor Research* 47:335-351.

KEDDY-HECTOR, D. 2000. Aplomado Falcon (*Falco femoralis*). In A. Poole [Ed.], The birds of North America online, No. 549. Cornell Lab of Ornithology, Ithaca, NY U.S.A. <http://bna.birds.cornell.edu/bna/species/549>, (last accessed 1 January 2015).

MACÍAS-DUARTE, A. 2002. Éxito reproductivo, presas potenciales y hábitat del halcón aplomado *Falco femoralis septentrionalis* Todd en Chihuahua, México. M.S. thesis, Universidad Autónoma de Chihuahua, Chihuahua, Chihuahua, Mexico.

———, A.B. MONTOYA, W.G. HUNT, A. LAFON-TERRAZAS, AND R. TAFANELLI. 2004. Reproduction, prey, and habitat of the Aplomado Falcon (*Falco femoralis*) in desert grasslands of Chihuahua, Mexico. *Auk* 121:1081-1093.

———, C.E. MÉNDEZ-GONZÁLEZ, J.R. RODRÍGUEZ-SALAZAR, W.G. HUNT, AND P.G. KRANNITZ. 2009. Factors influencing habitat use by migratory grassland birds in the state of Chihuahua, Mexico. *Auk* 126:896-905.

MEYER, R.A. AND S.O. WILLIAMS. 2005. Recent nesting and current status of the Aplomado Falcon (*Falco femoralis*) in New Mexico. *North American Birds* 2:352-356.

MONTOYA, A.B. 1995. Breeding biology of Aplomado Falcons in desert grasslands of Chihuahua, Mexico. M.S. thesis, New Mexico State University, Las Cruces, NM U.S.A.

———, P.J. ZWANK, AND M. CARDENAS. 1997. Breeding biology of Aplomado Falcons in desert grasslands of Chihuahua, Mexico. *Journal of Field Ornithology* 68:135-143.

POOL, D., A.O. PANJABI, A. MACÍAS-DUARTE, AND D.M. SOLHJEM. 2014. Rapid expansion of croplands in Chihuahua, Mexico threatens declining North American grassland bird species. *Biological Conservation* 170:274-281.

REYES-GOMEZ, V.M., S.C. DÍAZ, L. BRITO-CASTILLO, AND D. NÚÑEZ-LÓPEZ. 2013. ENSO/drought effects and their impact in the ecology and economy of the state of Chihuahua, Mexico. Pages 101-117 in A. Yañez, R. Dávalos-Sotelo, J.W. Day, and E. Reyes [Eds.], Ecological dimensions for sustainable socio economic development. WIT Press, Southampton, U.K.

WARNER, R.R. AND P.L. CHESSON. 1985. Coexistence mediated by recruitment fluctuations—a field guide to the storage effect. *American Naturalist* 125:769-787.

YOUNG, K.E., B.C. THOMPSON, A.L. TERRAZAS, A.B. MONTOYA, AND R. VALDEZ. 2004. Aplomado Falcon abundance and distribution in the northern Chihuahuan Desert of Mexico. *Journal of Raptor Research* 38:107-117.

Received 22 August 2015; accepted 17 November 2015
Associate Editor: Vincenzo Penteriani